

Quantifying the impact of future Sandage-Loeb test data on dark energy constraints

Jia-Jia Geng,^a Jing-Fei Zhang,^a Xin Zhang^{a,b,1}

^aDepartment of Physics, College of Sciences, Northeastern University,
Shenyang 110004, China

^bCenter for High Energy Physics, Peking University,
Beijing 100080, China

E-mail: gengjiajia163@163.com, jfzhang@mail.neu.edu.cn,
zhangxin@mail.neu.edu.cn

Abstract. The Sandage-Loeb (SL) test is a unique method to probe dark energy in the “redshift desert” of $2 \lesssim z \lesssim 5$, and thus it provides an important supplement to the other dark energy probes. Therefore, it is of great importance to quantify how the future SL test data impact on the dark energy constraints. To avoid the potential inconsistency in data, we use the best-fitting model based on the other geometric measurements as the fiducial model to produce 30 mock SL test data. The 10-yr, 20-yr, and 30-yr observations of SL test are analyzed and compared in detail. We show that compared to the current combined data of type Ia supernovae, baryon acoustic oscillation, cosmic microwave background, and Hubble constant, the 30-yr observation of SL test could improve the constraint on Ω_m by about 80% and the constraint on w by about 25%. Furthermore, the SL test can also improve the measurement of the possible direct interaction between dark energy and dark matter. We show that the SL test 30-yr data could improve the constraint on γ by about 30% and 10% for the $Q = \gamma H \rho_c$ and $Q = \gamma H \rho_{de}$ models, respectively.

¹Corresponding author.

Contents

1	Introduction	1
2	Methodology	2
3	Results and discussions	4
4	Summary	7

1 Introduction

Current cosmological observations, such as the type Ia supernovae (SN), the cosmic microwave background (CMB), the baryon acoustic oscillation (BAO), and so forth, have led to the precision cosmology, in which the geometry and matter/energy contents of the universe are precisely determined. However, the nature of dark energy still remains enigmatic. To determine the dark energy property, more highly accurate data are needed. In particular, it is of great importance to combine different observations to precisely determine the equation of state of dark energy. Different observational data are sensitive in different ways and degrees to dark energy parameters (as well as other parameters), and thus the alone use of one kind of data would lead to some parameter degeneracy, which might be broken when other kinds of observations are combined. In the forthcoming years, the close combination of geometric measurements, such as SN and BAO, and structure's growth measurements, such as clusters of galaxies counts and weak lensing, will definitely play a crucial role in determining the property of dark energy. On the other hand, some other promising observations will also provide important supplement, significantly affecting the constraints on dark energy.

The Sandage-Loeb (SL) test is a purely geometric measurement of the expansion of the universe, which was firstly proposed by Sandage [1] to directly measure the temporal variation of the redshift of extra-galactic sources, and then improved by Loeb [2] who provided a realistic way to measure the redshift drift by detecting the redshift variation of quasar (QSO) Lyman- α absorption lines. The SL test is rather important due to the fact that it is a unique method to explore the cosmic expansion (and, consequently, dark energy) at the “redshift desert” ($2 \lesssim z \lesssim 5$) that is not covered by any other dark energy probes. Combining the SL test data from this high-redshift region with other data from low-redshift region, such as SN and BAO, will definitely lead to significant impact on the dark energy constraints. The scheduled 39-meter Extremely Large Telescope (ELT) equipped with a high resolution spectrograph called the Cosmic Dynamics Experiment (CODEX) is designed to perform the SL test. Numerous works in this aspect have been done [3–9], some of which assume 240 observed QSOs in the simulations. However, according to an extensive Monte Carlo simulation [10], only about 30 QSOs will be bright enough and/or lie at a high enough redshift for observing redshift drift using a 39-m telescope. On the other hand, serious and careful quantification of the impact of future SL test data on dark energy constraints based on the observation of 30 QSOs is still absent. In this paper, we will provide such an analysis.

In existing papers on the SL test, the simulated SL test data are produced based on the fiducial (best-fitting) Λ CDM model, and constraints on other dark energy models are

performed in the light of these simulated data. Inconsistencies may occur when these simulated SL test data are combined with other actual data, and thus such a combination may not be appropriate. In order to properly quantify the impact of the future SL test data on dark energy constraints, producing the simulated SL test data consistent with other actual observations is particularly important and necessary. Here, we note that the SL test alone cannot provide a tight constraint on dark energy owing to the fact that its data are all within the high-redshift range ($2 \lesssim z \lesssim 5$) and the low-redshift data are lacking. Hence, the combination of SL test with low-redshift observations is a must for constraining dark energy. At present, in order to estimate the potential impact of the future SL test data on dark energy constraints, we have to combine the simulated SL test data with the currently available actual data covering the low-redshift region. And, to avoid the possible inconsistencies between the actual data and simulated future data, we will choose the best-fitting dark energy models using the current data as the fiducial models in producing the simulated future data. Such a scheme is of great importance in effectively quantifying how the future SL test data impact on dark energy constraints.

Typical dark energy models should be chosen as proper examples in the analysis. In this paper, we focus on the constant w dark energy model. Since dark energy along with cold dark matter (CDM) dominates the universe, we call this model the w CDM model for convenience. In addition, because there is always an important possibility that dark energy directly interacts with cold dark matter, we also consider the interacting extensions to the w CDM model. We call such interacting dark energy models the Iw CDM models.

In the cases of dark energy interacting with dark matter, the energy balance equations for dark energy and cold dark matter are

$$\dot{\rho}_{de} + 3H\rho_{de}(1+w) = -Q, \quad (1.1)$$

$$\dot{\rho}_c + 3H\rho_c = Q, \quad (1.2)$$

where the Hubble parameter $H = \dot{a}/a$ describes the expansion rate of the universe and the interacting term Q describes the energy transfer rate between dark energy and dark matter. The simplest cases are considered: (i) $Q = \gamma H\rho_c$, with which the model is called the Iw CDM1 model, and (ii) $Q = \gamma H\rho_{de}$, with which the model is called the Iw CDM2 model, where γ is the dimensionless coupling.

2 Methodology

For the current observations used in this analysis, we only consider the most typical ones. We consider a combination of SN, BAO, CMB, and H_0 , which is the most commonly used data combination in studies of dark energy. For the SN data, we use the SNLS compilation with a sample of 472 SNe [11]. For the BAO data, we use the $r_s/D_V(z)$ measurements from 6dFGS, SDSS-DR7, SDSS-DR9, and WiggleZ (six data from $z = 0.1$ to 0.73 , with the inverse of covariance matrix given in Ref. [12]). For the CMB data, we use the Planck distance priors given by Ref. [13]. Since we investigate the geometric measurements for dark energy constraints, the use of the CMB distance priors is fairly sufficient. We also use the direct measurement of H_0 by the cosmic distance ladder from the HST, $H_0 = 73.8 \pm 2.4 \text{ km s}^{-1} \text{ Mpc}^{-1}$ [14]. Dark energy models are firstly constrained by using the SN+BAO+CMB+ H_0 data combination, and the best-fitting models derived are then chosen as the fiducial models to produce the simulated SL test data. The resulting simulated SL test data are thus well

consistent with the actual current data, and consequently, the combination of the simulated SL test data with SN, BAO, CMB, and H_0 is appropriate.

Of course, in the 6-parameter base Λ CDM model, the Planck data are in tension with H_0 direct measurement at about the 2.5σ level, and are in tension with the SNLS data at about the 2σ level [15]. But, it is shown that the tension on H_0 could be greatly relieved once the dynamical dark energy is considered [16] (or a sterile neutrino species is included in the model [17–20]). The tension of Planck with SN may be due to some unknown systematic error sources in SN data, such as the absence of the consideration of possible evolution of the color-luminosity parameter β , as analyzed in Refs. [21, 22]. However, in this paper, we only focus on the impact of the SL test data, and so we omit the tiny inconsistency between Planck and SN.

As mentioned above, the SL test is a way to measure the redshift variation of quasar Lyman- α absorption lines. The velocity shift can be detected by subtracting the spectral templates of a quasar taken at two different times, and the quasar systems are now readily targeted and observable in the redshift range $2 \lesssim z \lesssim 5$. The redshift variation is defined as a spectroscopic velocity shift and expressed as [2]

$$\Delta v \equiv \frac{\Delta z}{1+z} = H_0 \Delta t_o \left[1 - \frac{E(z)}{1+z} \right], \quad (2.1)$$

where $E(z) = H(z)/H_0$ and Δt_o is the time interval of observation.

For the dark energy models considered in this paper, namely the w CDM, the Iw CDM1, and the Iw CDM2 models, the corresponding expressions of $E(z)$ are explicitly given in the following,

$$E(z)^2 = \Omega_r(1+z)^4 + \Omega_b(1+z)^3 + \Omega_c(1+z)^3 + (1 - \Omega_r - \Omega_b - \Omega_c)(1+z)^{3(1+w)}, \quad (2.2)$$

$$E(z)^2 = \Omega_r(1+z)^4 + \Omega_b(1+z)^3 + \frac{3w}{\gamma + 3w} \Omega_c(1+z)^{3-\gamma} + (1 - \Omega_r - \Omega_b - \frac{3w}{\gamma + 3w} \Omega_c)(1+z)^{3(1+w)}, \quad (2.3)$$

$$E(z)^2 = \Omega_r(1+z)^4 + \Omega_b(1+z)^3 + \frac{3w\Omega_c + \gamma(1 - \Omega_r - \Omega_b)}{\gamma + 3w} (1+z)^3 + \frac{3w(1 - \Omega_r - \Omega_b - \Omega_c)}{\gamma + 3w} (1+z)^{3(1+w)+\gamma}, \quad (2.4)$$

where Ω_r , Ω_b , and Ω_c are the present-day density parameters for the radiation, baryons, and cold dark matter, respectively, and $\Omega_r = \Omega_m/(1+z_{\text{eq}})$, with $\Omega_m = \Omega_c + \Omega_b$ and $z_{\text{eq}} = 2.5 \times 10^4 \Omega_m h^2 (T_{\text{cmb}}/2.7\text{K})^{-4}$ (here we take $T_{\text{cmb}} = 2.7255\text{ K}$).

Using Monte Carlo simulations and considering not just the Ly α forest but all available absorption lines, including metal lines, over the entire accessible optical wavelength range down to a QSO's Ly γ emission line, the accuracy of the spectroscopic velocity shift measurements expected by CODEX can be expressed as [10]

$$\sigma_{\Delta v} = 1.35 \left(\frac{S/N}{2370} \right)^{-1} \left(\frac{N_{\text{QSO}}}{30} \right)^{-1/2} \left(\frac{1+z_{\text{QSO}}}{5} \right)^f \text{ cm s}^{-1}, \quad (2.5)$$




Figure 1. Constraints on the w CDM, Iw CDM1, and Iw CDM2 models from the current SN+BAO+CMB+ H_0 data.

where S/N is the signal-to-noise ratio defined per $0.0125 \text{ } \ddot{\text{A}}$ pixel, N_{QSO} is the number of observed quasars, z_{QSO} represents their redshifts, and the exponent $f = -1.7$ for $2 < z < 4$ and $f = -0.9$ for $z > 4$. We choose $N_{\text{QSO}} = 30$ mock data uniformly distributed among the six redshift bins of $z_{\text{QSO}} \in [2, 5]$. The parameters in Eq. (2.1) are taken to be the best-fit values in the fit to the SN+BAO+CMB+ H_0 data, and error bars are computed from Eq. (2.5) with a $S/N = 3000$. We perform an MCMC likelihood analysis [23] to obtain $\mathcal{O}(10^6)$ samples for each model considered in this paper.

3 Results and discussions

Constraints on the w CDM (black), Iw CDM1 (purple), and Iw CDM2 (cyan) models from the current SN+BAO+CMB+ H_0 data are shown in Fig. 1. From the figure, one can clearly see that for all the three models $w < -1$ is preferred at more than 2σ level. The coupling γ is tightly constrained in the Iw CDM1 model, but its constraint is much weaker in the Iw CDM2 model. In both cases, $\gamma < 0$ is slightly favored, deviating from the noninteracting case ($\gamma = 0$) at about the 1.4σ level. (A $\gamma < 0$ indicates the energy transfer from dark matter to dark energy, in our convention.)




Figure 2. The SL test for the w CDM model. (a) Reconstructed velocity drift using the current SN+BAO+CMB+ H_0 constraint results for 10-yr, 20-yr, and 30-yr observations of SL test (red, green, and blue bands). Comparison is made with the error bars of the simulated SL test mock data. (b) Combined constraints on the w CDM model with current only, current+SL 10-yr, current+SL 20-yr, and current+SL 30-yr data. (c) Comparison of the constraints in the Ω_m - w plane from current only and SL test only data. Note that the orientations of the degeneracies of the two are nearly orthogonal, so the SL test is highly efficient in breaking the degeneracy in the other geometric measurements. In the SL test simulations, we assume $S/N = 3000$ and $N_{\text{QSO}} = 30$, where S/N is the signal-to-noise ratio defined per $0.0125 \text{ \AA}$ pixel and N_{QSO} is the number of observed quasars, and we also assume that the 30 mock data are uniformly distributed among the six redshift bins of $z_{\text{QSO}} \in [2, 5]$.

Simulating the 30 mock SL data using the best-fitting model as the fiducial model, and then combining them with the current real data, one can perform a joint constraint on the chosen dark energy model. The w CDM model is taken as a concrete example, and a detailed analysis is made for it. Figure 2 summarizes the results of our analysis.

In order to directly compare the accuracies of the current data and the future SL data, we reconstruct the velocity shifts for the w CDM model in the redshift range from $z = 0$ to 5.5 by using the fit results from the current SN+BAO+CMB+ H_0 data, plotted in Fig. 2(a) as colored bands. Red, green, and blue bands are for the 10-yr, 20-yr, and 30-yr velocity-shift reconstructions, respectively. The error bars in the SL test, estimated by Eq. (2.5), are also plotted in Fig. 2(a) on the corresponding bands, for a direct comparison. One can see that

the 10-yr observation of SL test is evidently worse than the current combined observations in accuracy, but a 20-yr observation can greatly improve the accuracy. The 30-yr observation of SL test gets further improvement, basically comparable with the current combined data in accuracy. So, the SL test, as a high-redshift supplement of $2 \lesssim z \lesssim 5$ to other cosmological geometric measurements, will play an important role in probing the expansion history of the universe and thus the property of dark energy.

Figure 2(b) shows a comparison of the joint constraints on the w CDM model in the Ω_m - w plane. The 68% and 95% CL posterior distribution contours are shown, where the current only, the current+SL 10-yr, the current+SL 20-yr, and the current+SL 30-yr results are colored with white, red, green, and blue, respectively. It is clear that using only the SL 10-yr combined data can improve the constraint significantly. Of particular importance is that the combination with the SL test can tightly constrain Ω_m to a high precision. With an only 10-yr observation, the constraint of Ω_m will be improved by 51.3%. That is an enormous improvement. The constraint on w can also be improved significantly. With an only 10-yr observation, the constraint of w is improved by 12.1%, which is also remarkable. With the 20-yr and 30-yr observations, the constraints on Ω_m are improved by 72.8% and 81.2%, respectively, and the constraints on w are improved by 18.1% and 25.0%, respectively. Therefore, we can conclude that using a 30-yr observation of the SL test the geometric constraints on dark energy would be improved enormously, i.e., the constraints on Ω_m and w would be improved, relative to the current joint observations, by at least 80% and 25%, respectively. Note that this is a conservative estimation since 30 years later other geometric measurements themselves would also be improved greatly in accuracy, and the combination of these highly accurate data could constrain dark energy more tightly.

In Fig. 2(c), we show how the SL test breaks the degeneracy in the current data constraints. Solely using the SL test data cannot place tight constraints on dark energy due to the lack of low-redshift data. In this case, strong degeneracy between w and Ω_m will appear, as shown in Fig. 2(c) where the 20-yr and 30-yr observations are taken as examples. However, one can also directly see that the orientations of the degeneracies for the SL test and the current combined data are nearly orthogonal, and so the SL test is highly efficient in breaking the degeneracy in the other geometric measurements. The 68% and 95% CL contours from the current+SL 20-yr (green) and current+SL 30-yr (blue) combined constraints are also plotted, from which one can directly see the crucial role played by the SL test in breaking the degeneracy. Further comparing the 20-yr and 30-yr observations of the SL test, we find that the 30-yr observation improves, relative to the 20-yr one, the constraints on Ω_m and w by 30.2% and 8.4%, respectively. Therefore, in the following, to show the decisional impact on dark energy constraints from the SL test, we use the 30-yr observation as an example.

At last, we discuss how the SL test helps improve the measurement of the possible direct interaction between dark energy and dark matter. In Fig. 3, we show the joint constraints on the two Iw CDM models in the Ω_m - w and the Ω_m - γ planes. The impact of the SL test can be directly seen from the comparison of the red (current only) and blue (current+SL 30-yr) contours. Compared to the current only data, the current+SL 30-yr data improve the constraints on Ω_m , w , and γ by 68.3%, 20.5%, and 29.5%, respectively, for the Iw CDM1 model, and by 58.0%, 14.7%, and 10.4%, respectively, for the Iw CDM2 model. Therefore, we find that as a purely geometric measurement, the SL test can also mildly improve the constraints on the interaction between dark energy and dark matter.




Figure 3. Constraints on the two $Iw\text{CDM}$ models from the current only and current+SL 30-yr data.

4 Summary

In this paper, we have discussed the quantification of the impact of future SL test data on dark energy constraints. The SL test is the unique dark energy probe of the “redshift desert” of $2 \lesssim z \lesssim 5$ and so it is particularly important, as a purely geometric measurement, for constraining dark energy as a supplement to other important geometric measurements. To quantify its impact, we should guarantee that the simulated future SL test data are consistent with the other geometric measurement data. Thus, we used the best-fitting models based on the other combined geometric measurement data as the fiducial models to produce the mock SL test data, and then do the analysis with these data. Typical dark energy models, such as the $w\text{CDM}$ model and its interacting extensions ($Iw\text{CDM}$ models), were considered in this analysis. Since we only focus on the purely geometric measurements, we do not consider the cosmological perturbations in the dark energy models. But we also note here that the cosmological perturbations would always be stable in all the interacting dark energy models, provided that a new framework for calculating the perturbations is adopted, as analyzed in detail in Ref. [24]. For related discussions concerning the stability, see also Ref. [25].

With the help of the SL test data, the constraints on dark energy will be improved significantly, compared to the current SN+BAO+CMB+ H_0 constraint results. The 10-yr,

20-yr, and 30-yr observations of SL test have been analyzed and compared in detail, and we found that the 30-yr observation could provide fairly important supplement to the other observations. The 30-yr observation of SL test is able to improve, relative to the current combined data, the constraint of Ω_m by about 80% and the constraint of w by about 25%, according to our analysis. In addition, for the Iw CDM model with $Q = \gamma H \rho_c$, the constraint on γ is improved by about 30%, and for the Iw CDM model with $Q = \gamma H \rho_{de}$, the constraint on γ is improved by about 10%. Therefore, according to the quantification study, the SL test is proven to be a very important supplement to the other geometric measurement observations, and will definitely play a crucial role in constraining dark energy in the future.

Acknowledgments

We acknowledge the use of **CosmoMC**. We thank Yun-He Li for helpful discussion. J.F.Z. is supported by the Provincial Department of Education of Liaoning under Grant No. L2012087. X.Z. is supported by the National Natural Science Foundation of China under Grant No. 11175042 and the Fundamental Funds for the Central Universities under Grant No. N120505003.

References

- [1] A. Sandage, “The Change of Redshift and Apparent Luminosity of Galaxies due to the Deceleration of Selected Expanding Universes,” *Astrophys. J.* **136**, 319 (1962).
- [2] A. Loeb, “Direct Measurement of Cosmological Parameters from the Cosmic Deceleration of Extragalactic Objects,” *Astrophys. J.* **499**, L111 (1998) [astro-ph/9802122].
- [3] P. -S. Corasaniti, D. Huterer and A. Melchiorri, “Exploring the Dark Energy Redshift Desert with the Sandage-Loeb Test,” *Phys. Rev. D* **75**, 062001 (2007) [astro-ph/0701433].
- [4] A. Balbi and C. Quercellini, “The time evolution of cosmological redshift as a test of dark energy,” *Mon. Not. Roy. Astron. Soc.* **382**, 1623 (2007) [arXiv:0704.2350 [astro-ph]].
- [5] H. -b. Zhang, W. Zhong, Z. H. Zhu and S. He, “Exploring holographic dark energy model with Sandage-Leob test,” *Phys. Rev. D* **76**, 123508 (2007) [arXiv:0705.4409 [astro-ph]].
- [6] J. Zhang, L. Zhang and X. Zhang, “Sandage-Loeb test for the new agegraphic and Ricci dark energy models,” *Phys. Lett. B* **691** (2010) 11 [arXiv:1006.1738 [astro-ph.CO]].
- [7] Z. Li, K. Liao, P. Wu, H. Yu and Z. -H. Zhu, “Probing modified gravity theories with the Sandage-Loeb test,” *Phys. Rev. D* **88**, 023003 (2013) [arXiv:1306.5932 [gr-qc]].
- [8] S. Yuan, S. Liu and T. -J. Zhang, “Breaking through the high redshift bottleneck of Observational Hubble parameter Data: The Sandage-Loeb signal Scheme,” arXiv:1311.1583 [astro-ph.CO].
- [9] M. -J. Zhang and W. -B. Liu, “Observational constraint on the interacting dark energy models including the Sandage-Loeb test,” arXiv:1312.0224 [astro-ph.CO].
- [10] J. Liske, A. Grazian, E. Vanzella, M. Dessauges, M. Viel, L. Pasquini, M. Haehnelt and S. Cristiani *et al.*, “Cosmic dynamics in the era of Extremely Large Telescopes,” *Mon. Not. Roy. Astron. Soc.* **386**, 1192 (2008) [arXiv:0802.1532 [astro-ph]].
- [11] A. Conley *et al.* [SNLS Collaboration], “Supernova Constraints and Systematic Uncertainties from the First 3 Years of the Supernova Legacy Survey,” *Astrophys. J. Suppl.* **192**, 1 (2011) [arXiv:1104.1443 [astro-ph.CO]].
- [12] G. Hinshaw *et al.* [WMAP Collaboration], “Nine-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Parameter Results,” *Astrophys. J. Suppl.* **208**, 19 (2013) [arXiv:1212.5226 [astro-ph.CO]].

- [13] Y. Wang and S. Wang, “Distance Priors from Planck and Dark Energy Constraints from Current Data,” *Phys. Rev. D* **88**, 043522 (2013) [arXiv:1304.4514 [astro-ph.CO]].
- [14] A. G. Riess *et al.*, “A 3% Solution: Determination of the Hubble Constant with the Hubble Space Telescope and Wide Field Camera 3,” *Astrophys. J.* **730**, 119 (2011) [arXiv:1103.2976 [astro-ph.CO]].
- [15] P. A. R. Ade *et al.* [Planck Collaboration], “Planck 2013 results. XVI. Cosmological parameters,” arXiv:1303.5076 [astro-ph.CO].
- [16] M. Li, X. -D. Li, Y. -Z. Ma, X. Zhang and Z. Zhang, “Planck Constraints on Holographic Dark Energy,” *JCAP* **1309**, 021 (2013) [arXiv:1305.5302 [astro-ph.CO]].
- [17] J. -F. Zhang, Y. -H. Li and X. Zhang, “Sterile neutrinos help reconcile the observational results of primordial gravitational waves from Planck and BICEP2,” arXiv:1403.7028 [astro-ph.CO].
- [18] C. Dvorkin, M. Wyman, D. H. Rudd and W. Hu, “Neutrinos help reconcile Planck measurements with both Early and Local Universe,” arXiv:1403.8049 [astro-ph.CO].
- [19] J. -F. Zhang, Y. -H. Li and X. Zhang, “Cosmological constraints on neutrinos after BICEP2,” arXiv:1404.3598 [astro-ph.CO].
- [20] Y. -H. Li, J. -F. Zhang and X. Zhang, “Tilt of primordial gravitational wave spectrum in a universe with sterile neutrinos,” *Sci. China Phys. Mech. Astron.* **57**, 1455 (2014) [arXiv:1405.0570 [astro-ph.CO]].
- [21] S. Wang, Y. -H. Li and X. Zhang, “Exploring the evolution of color-luminosity parameter β and its effects on parameter estimation,” *Phys. Rev. D* **89**, 063524 (2014) [arXiv:1310.6109 [astro-ph.CO]].
- [22] S. Wang, J. -J. Geng, Y. -L. Hu and X. Zhang, “Revisit of constraints on holographic dark energy: SNLS3 dataset with the effects of varying β and different light-curve fitters,” arXiv:1312.0184 [astro-ph.CO].
- [23] A. Lewis and S. Bridle, “Cosmological parameters from CMB and other data: a Monte-Carlo approach,” *Phys. Rev. D* **66**, 103511 (2002) [astro-ph/0205436].
- [24] Y. -H. Li, J. -F. Zhang and X. Zhang, “New framework for calculating cosmological perturbations in a scenario of dark energy interacting with cold dark matter,” arXiv:1404.5220 [astro-ph.CO].
- [25] Y. -H. Li and X. Zhang, “Large-scale stable interacting dark energy model: Cosmological perturbations and observational constraints,” *Phys. Rev. D* **89**, 083009 (2014) [arXiv:1312.6328 [astro-ph.CO]].